

# SUPERDEFORMATION IN $^{147,148}\text{Eu}$ : IDENTICAL BANDS AND $\pi6^1 - \pi6^3$ CROSSINGS<sup>1</sup>

D. S. Haslip<sup>2</sup>, N. Kintz<sup>3</sup>, S. Flibotte<sup>2</sup>, R. A. E. Austin<sup>2</sup>, G. de France<sup>4</sup>, M. Devlin<sup>5</sup>, Ch. Finck<sup>3</sup>, A. Galindo-Uribarri, G. Gervais<sup>2</sup>, D. R. LaFosse<sup>5</sup>, T. J. Lampman<sup>2</sup>, I. Y. Lee<sup>6</sup>, F. Lerma<sup>5</sup>, A. O. Macchiavelli<sup>6</sup>, R. W. MacLeod<sup>6</sup>, S. M. Mullins<sup>7</sup>, J. M. Nieminen<sup>2</sup>, T. Rodinger<sup>2</sup>, D. G. Sarantites<sup>5</sup>, O. Stezowski<sup>3</sup>, C. E. Svensson<sup>2</sup>, J. P. Vivien<sup>3</sup>, J. C. Waddington<sup>2</sup>, D. Ward<sup>6</sup>, and J. N. Wilson<sup>2</sup>

Eight superdeformed bands have been observed in the nuclei  $^{147,148}\text{Eu}$ . Some of these are identical to bands in the Z+1 nuclei  $^{148,149}\text{Gd}$ , and are interpreted as having the same single-particle configurations with the exception of a hole in one of the [301]1/2 proton orbitals. Some of the bands in  $^{147}\text{Eu}$  exhibit either upbends or downbends in their moments of inertia which are related to the crossing of  $\pi6^1$  and  $\pi6^3$  proton intruder configurations. These results constitute the first evidence of this crossing.

---

<sup>1</sup>Abstract of published paper: Phys. Rev. C **57**, 2196 (1998).

<sup>2</sup>McMaster University, Hamilton, Ontario, L8S 4M1 Canada.

<sup>3</sup>Institut de Recherches Subatomiques, F-67037 Strasbourg, France.

<sup>4</sup>GANIL, B.P. 5027, F-4076 Caen, France.

<sup>5</sup>Washington University, St. Louis, MO 63130.

<sup>6</sup>Lawrence Berkeley National Laboratory, Berkeley, CA 94720.

<sup>7</sup>RSPHYSSE, ANU, Canberra, ACT 0200, Australia.